



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.


III. Solution by the PROPOSER.

The roots of $ax^2 + 2bx + c = 0$ and $a'x^2 + 2b'x + c' = 0$ will be harmonic if $ac' + a'c - 2bb' = 0$ (see Scott's Geometry, page 45).

Let $x^2 = p^2$ give the points A and B . Let $x = OM = K < (OB = p)$ be midway between the other points, P and Q . The equation giving P and Q is

$$a'x^2 + 2b'x + c' = 0, \text{ with the conditions } \frac{b'}{a} = -K, \text{ and } c' - p^2 a' = 0,$$

$$\text{or } x^2 - 2Kx + p^2 = 0.$$



But since $K < p$, $K^2 - p^2 < 0$, the roots of this equation are imaginary, and since there are an indefinite number of values for $K < p$, there will be an indefinite number of pairs of imaginary points on the line harmonic with the given real pair. (Scott's Geometry, page 45.)

Solved in a similar manner by *G. B. M. ZERR*.

PROBLEMS.

63. Proposed by *ALFRED HUME*, C. E., D. Sc., Professor of Mathematics, University of Mississippi, P. O. University, Mississippi.

Prove, analytically :—A rectangular hyperbola cannot be cut from a right circular cone unless the angle at its vertex is greater than a right angle.

64. Proposed by *WILLIAM E. HEAL*, Member of the London Mathematical Society and Treasurer of Grant County, Marion, Indiana.

Let the bisectors of the angles A, B, C of a triangle meet the sides opposite A, B, C in A', B', C' . Let AA', BB', CC' meet the sides of the triangle $A'B'C'$ in A'', B'', C'' . Let this process continue indefinitely. Express the sides and angles of the triangle $A^{(m)}B^{(m)}C^{(m)}$ in terms of the sides and angles of the original triangle ABC .